

Katalog von 32 „Neuen“ Sternen.

Nr.	Name	Ort für 1900.0		Präzession 1900		Kartenort		Größe			Entdecker	Jahr
		AR.	Dekl.	AR.	Dekl.	AR.	Dekl.	Max.	Min.			
1	Nova B Cassiopejæ	0 ^h 19 ^m 15 ^s	+63° 35' 5	+3.30	+0.33	0 ^h 16 ^m 47 ^s	+63° 20.6	> 1 ^m	10 ^m		Tycho Brahe	1572
2	» Piscium	0 29 49	+ 9 45	+3.10	+0.33	0 27 18	+ 9 30	8.8	<14	ph	Ernst	1907
3	» S Andromedæ	0 37 15	+40 43.2	+3.26	+0.33	0 34 49	+40 28.3	5	<12		Hartwig	1885
4	» V Persei 1	1 55 6	+56 15	+4.00	+0.29	1 52 6	+56 2	9.3	<15	ph	Fleming	1887
5	» Arietis 1	2 42 56	+16 56.9	+3.34	+0.25	2 40 26	+16 45.4	9.5	—		Krueger	1854
6	» Arietis 2	3 19 19	+19 29.6	+3.43	+0.22	3 16 45	+19 19.9	<11	13.5	ph	M. Wolf	1905
7	» Persei 2	3 24 24	+43 33.7	+4.06	+0.21	3 21 22	+43 24.2	0	12		Anderson	1901
8	» T Aurigæ	5 25 34	+30 22.2	+3.85	+0.05	5 22 41	+30 19.9	4.5	<15		Anderson	1892
9	» Geminorum 1	6 37 49	+30 2.7	+3.83	-0.05	6 34 56	+30 5.0	7.3	12.5	ph	Turner	1903
10	» Geminorum 2	6 48 25	+32 16.0	+3.89	-0.07	6 45 30	+32 19.0	4	<11		Enebo	1912
11	» Velorum	10 58 20	-53 50.9	+2.59	-0.32	10 57 15	-53 42.9	9.7	<15	ph	Leavitt	1906
12	» RS Carinae	11 3 54	-61 23.6	+2.48	-0.32	11 2 52	-61 15.6	8	13	ph	Fleming	1895
13	» Z Centauri	13 34 17	-31 7.6	+3.40	-0.30	13 32 52	-31 0.0	7.2	<11.5	ph	Fleming	1895
14	» Circini	14 40 31	-59 35.0	+4.54	-0.26	14 38 38	-59 29.0	9.5	15	ph	Leavitt	1907
15	» Normæ	15 22 11	-50 13.9	+4.31	-0.21	15 20 23	-50 8.6	9.6	13	ph	Fleming	1893
16	» T Coronæ	15 55 19	+26 12.2	+2.51	-0.17	15 53 26	+26 20.1	2.0	9.5		Birmingham	1866
17	» T Scorpii	16 11 5	-22 43.6	+3.57	-0.15	16 8 25	-22 36.7	7.0	<12		Auwers	1860
18	» Arae	16 33 1	-52 13.5	+4.67	-0.12	16 31 4	-52 10.4	6.0	10.0		Fleming	1910
19	» Ophiuchi 2	16 53 54	-12 44.4	+3.36	-0.10	16 51 23	-12 40.0	5.5	12.5		Hind	1848
20	» Ophiuchi 1	17 24 38	-21 23.7	+3.59	-0.05	17 21 57	-21 21.2	> 1	9.0		Brunowsky	1604
21	» Scorpii 2	17 47 28	-34 19.7	+3.98	-0.02	17 45 48	-34 19.2	8.8	—		Cannon	1906
22	» Sagittarii 2	17 53 49	-27 32.8	+3.77	-0.01	17 52 14	-27 32.5	7.8	10.5	ph	Fleming	1910
23	» Sagittarii 4	18 0 27	-27 26.5	+3.77	0.00	17 58 53	-27 26.5	10.4	<16	ph	Cannon	1901
24	» Sagittarii 3	18 13 48	-25 13.5	+3.70	+0.02	18 12 15	-25 14.0	8.5	<13		Cannon	1899
25	» Sagittarii 1	18 56 13	-13 18.2	+3.38	+0.08	18 53 41	-13 21.8	4.7	<14		Fleming	1898
26	» Aquilæ 2	18 56 48	- 4 35.2	+3.18	+0.08	18 54 25	- 4 38.8	6.5	—		Fleming	1905
27	» Aquilæ 1	19 15 16	- 0 19.2	+3.08	+0.11	19 12 58	- 0 24.2	7	<13		Fleming	1899
28	» Sagittæ	19 28 16	+17 32.0	+2.68	+0.13	19 26 15	+17 26.4	6	<11		d'Agelet	1783
29	» II Vulpeculæ	19 43 28	+27 4.2	+2.46	+0.15	19 41 37	+26 57.7	3	< 6		Anthelme	1670
30	» P Cygni 1	20 14 6	+37 43.3	+2.21	+0.18	20 12 27	+37 35.1	3.5	< 6		W. J. Blæu	1600
31	» Q Cygni 2	21 37 47	+42 23.1	+2.36	+0.27	21 36 1	+42 11.0	3	14.8		Schmidt	1876
32	» Lacertæ	22 31 45	+52 11.9	+2.42	+0.31	22 29 56	+51 58.0	8.0	13.1		Espin	1910